

- steroid with the "unnatural" 14 β configuration from the marine sponge *Petrosia Contignata* Thiele 1899 [J]. J Org Chem, 1992, 57: 525.
- [7] Bramley AM, Langlands JM, Burgoyne DL. IZP-94005, Contignastero-Pneumocort RTM [J]. Drugs of the Future, 1994, 19(8): 738.
- [8] Shen YP and Burgoyne DL. Efficient synthesis of IPL576, 092: A novel Antithrombotic Agent [J]. J Org Chem, 2002, 67: 3908.
- [9] Fabrizio DP, Agostino C, et al. molecular basis of phospholipase A2 inhibition by Petrospongiolide M [J]. Chem Bio Chem, 2002, 3: 664.
- [10] Potts BCM, Faulkner DJ, Jacobs RS. Phospholipase A₂ inhibitors from marine organisms [J]. J Nat Prod, 1992, 55: 1701.
- [11] Wen SJ, Hu TS, Yao ZJ. Macrocyclization studies and total synthesis of cyclomarin C, an anti-inflammatory marine cyclopeptide [J]. Tetrahedron, 2005, 61: 4931.
- [12] Antonio R, Giuseppe B, Clelia G. Halipeptins A and B: Two novel potent anti-inflammatory cyclic depsipeptides from the Vanuatu marine sponge *Haliclona* species [J]. J Am Chem Soc, 2001, 123: 10870.
- [13] Souuke H, Kazuishi M, Yasumasa H. Total synthesis of halir peptin A, a potent anti-inflammatory cycloepsipeptide from a marine sponge [J]. Tetrahedron Letters, 2006, 47: 1081.
- [14] Jiang ZD, Paul R. Lobophorins A and B, new anti-inflammatory macrolides produced by a tropical marine bacterium [J]. Biorganic & Medicinal Chemistry Letters, 1999, 9: 2003.
- [15] Jin DQ, Lin CS, Sung JY, et al. *Ulva conglobata* amarine algae, has neuroprotective and anti-inflammatory effects in murine hippocampal and microglial cells [J]. Neuroscience Letters, 2006, 402: 154.
- [16] McGee EG, McGee PL. Inflammatory processes in Alzheimer's disease [J]. Prog Neuropsychopharmacol Biol Psychiatry, 2003, 27: 741.
- [17] Jin WY, Se YY, Soo JO, et al. Bifunctional effects of fucoidan on the expression of inducible nitric oxide synthase [J]. Biophysical and Biophysical Research Communications, 2006, 346: 345.
- [18] Szabo C. Role of nitric oxide in endotoxic shock: An overview of recent advances [J]. Ann NY Acad Sci, 1998, 851: 422.
- [19] Weinberg JB. Nitric oxide synthase 2 and cyclooxygenase 2 interactions in inflammation [J]. Immunol Rev, 2000, 22: 319.

收稿日期: 2007-03-10

• 药物不良反应个案报告 •

苦碟子注射液致病人头痛眩晕 1例

蒋 李, 曹洪江, 高 伟(中国人民解放军第 456 医院药剂科, 山东 济南 250031)

中图分类号: R286

文献标识码: D

文章编号: 1006-0111(2008)02-0096-01

1 临床资料

患者, 男, 54岁, 住院号 141536 因冠心病于2007年6月15日来我院心血管内科就诊, 医嘱给予苦碟子注射液20mL, qid, 舒血宁注射液10mL, qid。首次给药给予0.9%氯化钠注射液+苦碟子注射液(沈阳双鼎制药有限公司, 批号: 06110101)20mL静脉滴注, 10min后患者出现轻微头痛、眩晕, 疑为滴速过快, 请示医生后减慢滴速, 15min后患者表述仍有眩晕感, 且不能耐受, 立即停药, 并给予吸氧治疗, 20min后患者眩晕感基本消失, 次日再次应用苦碟子注射液20mL静滴时, 又出现头痛、眩晕, 考虑为苦碟子注射液引起, 即停止苦碟子注射液治疗。

2 讨论

苦碟子注射液是由菊科植物苦碟子(学名抱茎苦卖菜)制成的静脉注射液, 主要成分为腺苷类及黄酮类物质。苦碟子具有扩张脑血管, 改善循环、增强纤溶酶活性, 抑制血小板聚集, 消除氧自由基等药理作用, 广泛用于脑梗死、冠心病的治疗。药物说明书未有有关不良反应的说明, 文献报道亦少见。本例患者为冠心病病人, 入院后应用苦碟子注射液后出现头痛、眩晕。故考虑为苦碟子注射液引起, 随即停用该药, 上述症状消失; 1d后重复应用该药, 又发生上述情况, 停药后消失。病人头痛、眩晕与应用苦碟子注射液密切相关, 其原因有待进一步研究, 提醒应用时注意。

收稿日期: 2007-07-16